

Protective effect of the hooks and stems of *Uncaria sinensis* against nitric oxide donor-induced neuronal death in cultured cerebellar granule cells

Yutaka SHIMADA,^{*a)} Koichi YOKOYAMA,^{a)} Hirozo GOTO,^{b)} Iwao SAKAKIBARA,^{c)} Nobuyasu SEKIYA,^{a)}
Naoki MANTANI,^{a)} Shinya SAKAI,^{a)} and Katsutoshi TERASAWA^{a)}

^{a)}Department of Japanese Oriental Medicine, Faculty of Medicine,

^{b)}Department of Kampo Diagnostics, Institute of Natural Medicine, Toyama Medical and Pharmaceutical University,
2630 Sugitani, Toyama, Japan.

^{c)}Tsumura Central Research Laboratories, 3586 Yoshiwara, Amimachi, Ibaraki 300-1192, Japan.

(Received January 7, 2002. Accepted February 4, 2002.)

Abstract

We have previously shown that an aqueous extract of the hooks and stems of *Uncaria sinensis* (OLIV.) HAVIL., *Uncariae Uncus Cum Ramulus*, protects against glutamate-induced neuronal death *in vitro*. Nitric oxide (NO) free radicals are also implicated in the process of neuronal death. In this study, we investigated the protective effects of *Uncaria sinensis* extract (USE) and its phenolic and alkaloid fractions against NO donors, sodium nitroprusside (SNP) and 3-morpholinosydnonimine (SIN-1), -induced neuronal death in cultured rat cerebellar granule cells. MTT assay showed cell viability to be significantly increased by the addition of USE (10, 30 and 100 $\mu\text{g/ml}$) compared with exposure (6, 12 and 24 h) to SNP (30 μM) only, and by the addition of USE (10 and 30 $\mu\text{g/ml}$) compared with exposure (6, 12 and 24 h) to SIN-1 (300 μM) only. Phenolic fraction of USE (10 and 30 $\mu\text{g/ml}$) significantly protected against SNP (30 μM , 24 hr)-induced cell death, and 3 and 10 $\mu\text{g/ml}$ of this fraction significantly protected against SIN-1 (300 μM , 24 hr)-induced cell death. Alkaloid fraction of USE (30 and 100 $\mu\text{g/ml}$) significantly protected against SNP (30 μM , 24 hr) and SIN-1 (300 μM , 24 hr)-induced cell death. These results appear to indicate that *Uncaria sinensis* has a protective effect against NO-mediated neuronal death in cultured cerebellar granule cells and that its active components are included in phenolic and alkaloid fractions.

Key words *Uncaria sinensis*, phenol, alkaloid, nitric oxide donor, neuronal death, protective effect.

Abbreviations CGC, cerebellar granule cell; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium bromide; NO, nitric oxide; SIN-1, 3-morpholinosydnonimine; SNP, sodium nitroprusside; US, *Uncaria sinensis*; USE, *Uncaria sinensis* extract.

Introduction

Nitric oxide (NO) plays an important role in the central nervous system as a messenger molecule. However, excessive generation of NO has been implicated in neurotoxicity. This form of neurotoxicity is thought to be concerned with a final common pathway of injury in a wide variety of neurological disorders including ischemia.¹⁾ During brain ischemia, excessive amounts of glutamate are released from the pre-synaptic site of

glutamate neuron, overstimulate glutamate receptors, especially *N*-methyl-D-aspartate (NMDA) receptors of neurons, finally leading to a series of potentially neurotoxic events through excessive Ca^{2+} influx.^{2,3)} One of these events is the activation of the neuronal form of NO synthase (nNOS) and the subsequent production of NO free radicals (NO^{\cdot}).^{1,2,4)} Moreover, some kinds of cytokines overstimulate astroglia or microglia after the brain ischemia/reperfusion and this leads to excessive and uncontrolled production of NO^{\cdot} through a Ca^{2+} -independent and inducible form of NO synthase

*To whom correspondence should be addressed. e-mail : shimada@ms.toyama-mpu.ac.jp

(iNOS).^{1,5} Another event is the generation of superoxide anion ($O_2^{\cdot-}$) due in part to the release of arachidonic acid.⁶ NO^{\cdot} can react with $O_2^{\cdot-}$ to form peroxynitrate ($ONOO^-$), which results in dose-dependent neuronal cell death.⁷ The mechanisms proposed for NO-mediated neurotoxicity include inactivation of the mitochondrial respiratory chain, nitrosylation of glyceraldehyde-3-phosphate dehydrogenase (GAPDH), inhibition of cis-aconitase, activation of poly (adenosine 5'-diphosphoribose (ADP-ribose)) synthetase, and DNA damage.^{1,8} Many investigators have reported that the application of various kinds of NO donors can produce cell death in cultured neurons.^{4,7-10} Sodium nitroprusside (SNP) generates NO^{\cdot} that reacts with endogenous $O_2^{\cdot-}$, 3-morpholinopyridone (SIN-1) has been shown to produce NO^{\cdot} and $O_2^{\cdot-}$, thereby producing peroxynitrate ($ONOO^-$) further hydroxyl radicals ($\cdot OH$).^{4,11}

The hooks and stems of *Uncaria sinensis* (OLIV.) HAVIL. (US), *Uncariae Uncus Cum Ramulus*, comprise the main medicinal plant of Choto-san (Diao-Teng-San in Chinese). Choto-san, which is composed of 11 crude drugs, has long been administered as a decoction to relatively aged patients suffering from headache, dizziness, vertigo, tinnitus, shoulder stiffness and so forth in China and Japan. We recently revealed the effectiveness of Choto-san on patients with vascular dementia by a well-controlled study and a double-blind, placebo-controlled study.^{12,13} We also demonstrated that the *Uncaria sinensis* water extract (USE) and its phenolic and alkaloid compounds have protective effects against glutamate-induced neuronal death in cultured cerebellar granule cells (CGCs).¹⁴⁻¹⁶ However, the effect of US against NO-mediated neurotoxicity has not been clarified. In this study, we investigated the neuroprotective effect of US from the view point of NO-mediated toxicity and found that US and its phenolic and alkaloid fractions have protective abilities against NO donor-induced neuronal death in CGCs.

Materials and Methods

Plant material and extraction: The US used in this study was obtained as described previously.¹⁷ It was purchased from a Chinese market in the Guangxi district in 1996, and voucher specimens (No. 16130) have been deposited in the herbarium stock room of the Tsumura

Laboratory.

Plant materials (1 kg) were chopped and extracted with boiling water (5 l) for 2 h three times. Each extract was combined and lyophilized to give a brown mass (114.6 g, yield 11.5%). The water extract was chromatographed on polus polymer gel (Diaion HP-20, 2 l), eluted with water (2 l, 58.2 g, yield 5.8 %; water eluate (USE-W)), H_2O -methanol (1:1) (2 l, 29.4 g, yield 2.9 %; phenolic fraction (USE-P)), and then methanol (2 l, 2.4 g, yield 0.2 %; alkaloid fraction (USE-A)), successively.¹⁵⁻¹⁷

CGCs culture: Cerebellar granule cells from 8-day-old Wistar rats were cultured in poly-L-lysine (Sigma) coated 35-mm culture dishes with basal Eagle medium (Sigma) containing 10% fetal bovine serum (Sigma) and 25 mM KCl at a density of 6×10^5 cells/ml (2 ml/dish) as described previously.¹⁴⁻¹⁶ The cultures were maintained at 37°C with 5% CO_2 in a humidified incubator, and cytosine arabinoside (Sigma) (10 μM) was added 18 h after plating to prevent proliferation of glial cells. The culture medium was not changed thereafter. The cells were used at 8-9 days *in vitro* for the experiments. All animal use procedures were approved by the Committee on Animal Experimentation at Toyama Medical and Pharmaceutical University.

Drug treatment and evaluation of cell viability: SNP, SIN-1 and 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) were purchased from Sigma. USE, USE-W, USE-P and USE-A were dissolved in Locke's solution (in mM: 154 NaCl, 5.6 KCl, 1.2 Mg, 3.6 $NaHCO_3$, 5.0 HEPES, 2.3 $CaCl_2$, 5.6 glucose, pH 7.4) as $\times 100$ concentrated stock solution, and added (20 μl) to the original culture medium. Only the same amount of Locke's solution was added to untreated and NO donor only (control) culture medium. Ten minutes later, the NO donors, SNP and SIN-1, were dissolved in Locke's solution as $\times 100$ concentrated solution and added (20 μl) to culture medium. The same amount of Locke's solution was added to only untreated cultures. The cell cultures were incubated with drugs at 37°C with 5% CO_2 in a humidified incubator for certain periods. Cell viability was evaluated by MTT assay as previously described.¹⁴⁻¹⁶ After incubation with drugs, MTT (500 $\mu g/ml$) was applied and incubated for 30 min at 37°C. Cells were then washed and lysed in isopropanol with 0.04 N HCl to dissolve the blue formazan products. Optical density was read at 570 nm with a spectropho-

tometer and expressed as percentage of the untreated cultures.

Statistical analysis : Values were expressed as mean \pm S.D. from 4 experiments. The data were analyzed by Kruskal-Wallis test followed by Bonferroni's *t*-test. A *p* value < 0.05 was considered statistically significant.

Results

Concentration relationship of NO donor-induced cell death and protective effect of USE

To establish a model of NO donor-induced neurotoxicity and examine the effect of USE in CGCs, we exam-




Fig. 1 Effect of *Uncaria sinensis* extract (USE) against SNP (a) and SIN-1 (b) -induced neuronal death evaluated by MTT assay in cultured cerebellar granule cells. USE was added to the culture medium 10 min before NO donors application. Cells were incubated in drugs for 24 h. Values are mean \pm S.D. from 4 experiments. **p*<0.05, ***p*<0.01 compared with control.




Fig. 2 Phase-contrast photographs of cerebellar granule cells. SNP (30 μM, 24 h) (a) and SIN-1 (300 μM, 24 h) (b) -induced neuronal degeneration appeared in the form of destruction of the soma and neurite network, and a bright pyknotic mass of nuclei. These toxicities were prevented by the application of *Uncaria sinensis* extract (USE) (c SNP 30 μM, USE 100 μg/ml; d SIN-1 300 μM, USE 30 μg/ml). Bar indicates 50 μm.

ined the concentration-response relationship of neuronal cell death at 24 h after exposure to NO donors. SNP (10, 30 and 100 μM) decreased cell viability dose-dependently (10 μM, 53 %; 30 μM, 15 %; 100 μM, 11 %) (Fig. 1a). SIN-1 (300 μM) also decreased cell viability (35 %) (Fig. 1b). USE had significant protective effects against SNP (10 and 30 μM)-induced cell death at concentrations of 10, 30 and 100 μg/ml, and against SNP (100 μM)-induced cell death at concentrations of 30 and 100 μg/ml, in a dose-dependent manner (Fig. 1a). Furthermore, USE (10 and 30 μg/ml) significantly prevented SIN-1 (300 μM)-induced cell death in a dose-dependent manner (Fig. 1b). SNP or SIN-1-induced neuronal degeneration appeared as destruction of the soma and neurite network as well as a bright pyknotic mass of nuclei (Fig. 2a,b). These indications of neuronal death were prevented by USE (Fig. 2c,d).

Time course of NO donor-induced cell death and protective effect of USE

Next, we examined the time course of NO donor-induced cell death and the protective effect of USE. Based on the results from the concentration relationship of SNP and SIN-1-induced cell death, we decided on concentrations of 30 μM of SNP and 300 μM of SIN-1. Application of SNP or SIN-1 for 3 h did not influence cell viability (SNP, 90 %; SIN-1, 99 %), but longer applications decreased cell viability (SNP 6 h, 44 %; 12 h, 33 %; 24 h, 24 %; SIN-1 6 h, 61 %; 12 h, 33 %; 24 h, 29 %) (Fig. 3a,b). USE (10, 30 and 100 μg/ml) signifi-




Fig. 3 Time course of SNP (30 μM) and SIN-1 (300 μM) -induced neuronal death and protective effect of *Uncaria sinensis* extract (USE) evaluated by MTT assay in cultured cerebellar granule cells. USE was added to the culture medium 10 min before NO donors application. Cells were incubated in drugs for 3, 6, 12 or 24 h. Values are mean \pm S.D. from 4 experiments. **p*<0.05, ***p*<0.01 compared with control at each time point.




Fig. 4 Effects of water eluate (USE-W), phenolic fraction (USE-P) and alkaloid fraction (USE-A) of *Uncaria sinensis* extract against SNP (30 μM, 24 h) (a) and SIN-1 (300 μM, 24 h) (b) -induced neuronal death evaluated by MTT assay in cultured cerebellar granule cells. Each fraction was added to the culture medium 10 min before NO donors application. Cells were incubated in drugs for 24 h. Values are mean ± S.D. from 4 experiments. * $p < 0.05$, ** $p < 0.01$ compared with control.

cantly prevented SNP-induced cell death at 6, 12 and 24 h (Fig. 3a). Furthermore, USE (10 and 30 μg/ml) provided significant protection against neuronal death induced by SIN-1 at 6, 12 and 24 h (Fig. 3b).

Protective effects of phenolic and alkaloid fractions against NO donor-induced cell death

We examined the protective effects of USE-W, USE-P and USE-A against SNP and SIN-1-induced neuronal death. USE-W had no protective ability (Fig. 4a,b). USE-P (10 and 30 μg/ml) and USE-A (30 and 100 μg/ml) provided protection against neuronal death induced by SNP (30 μM, 24 h) in a dose-dependent manner (Fig. 4a). USE-P (3 and 10 μg/ml) and USE-A (30 and 100 μg/ml) also had protective effects against neuronal death induced by SIN-1 (300 μM, 24 h) in a dose-dependent manner (Fig. 4b).

Finally, in order to examine whether USE and its fractions were toxic to CGCs, we evaluated the effects of them on cell viability in the absence of NO donors. As a result, USE and its fractions (24 h-incubation) did not show any toxicity at least at the concentrations of 100 μg/ml or less (data not shown).

Discussion

In the present study, we showed that USE has a neuroprotective effect against NO donor-induced cell death in CGCs and that its active components were included in phenolic and alkaloid fractions. Previously, we

reported that USE and its phenolic and alkaloid compounds protected against glutamate-induced neuronal death in CGCs.¹⁴⁻¹⁶ It is known that NO mediates glutamate neurotoxicity.^{1,4,10} The excessive amounts of glutamate, which are released from the pre-synaptic neurons during brain ischemia, overstimulate glutamate receptors in the post-synaptic neurons, especially NMDA receptors. This event leads to the excessive Ca²⁺ influx into neurons through voltage-dependent and more glutamate-regulated Ca²⁺-channels, then the activation of nNOS and subsequent production of NO.¹⁻⁴ Therefore the present and our previous studies suggest a possibility that the protective effect of USE against glutamate-induced neuronal death is at least in part due to its protective ability against NO-mediated neurotoxicity.

Glial cells also play important roles in the process of the neuronal death induced by brain ischemia. Glial cells take in glutamate and metabolize it to glutamine by glutamine synthetase.¹⁸ This ability is thought to act neuroprotectively in brain ischemia. On the other hand, in the later phase of brain ischemia/reperfusion, several stimuli involving activation of some kinds of cytokines, strongly induce iNOS expression in glial cells, allowing excessive and uncontrolled production of NO, and leading to delayed neuronal death.^{1,5} In cultured neurons, neuronal cell death produced by a low level of NO donor is characterized by delayed neuronal death, which is apoptosis rather than necrosis.⁷ In our present study, application of NO donors did not influence cell viability until 3 h, but longer application decreased cell viability and produced nuclear condensation and fragmentation in CGCs, a feature of apoptosis, and these findings were protected by USE. Whereas we prevented the proliferation of glial cells by application of cytosine arabinoside in the present cultures, further studies need to focus on whether US has any effect on glial cells in the process of neuronal death.

In the process of brain ischemia/reperfusion, large amounts of O₂⁻ are generated resulting from the mitochondrial dysfunction, the accumulation of hypoxanthine, the release of arachidonic acid, the activation of leukocytes, and so forth.^{1,6} Increasing evidences obtained from *in vivo* studies suggest that formation of ONOO⁻ from the reaction of NO with O₂⁻, and further OH generation, is a mechanism through which neurotoxicity is induced.^{1,7} In the present study, we used

two kinds of NO donors, SNP and SIN-1. At least on condition that SNP (30 μ M, 24 hr) and SIN-1 (300 μ M, 24 hr) were applied, phenolic fraction of US showed a protective effect at relatively lower concentration compared with alkaloid fraction.

Uncaria genus is known to have antioxidant and free radical scavenging activities from evaluation using the electron spin resonance technique and the inhibitory effect on lipid peroxidation in the brain of iron-induced epileptic rats.¹⁹⁾ Further, Uncaria genus has an inhibitory effect against convulsion induced by the *in vivo* administration of excitatory amino acids, glutamate and kainate,^{20,21)} the neuronal damages of which are thought to be caused in part by oxidative stress produced by NO and O₂⁻ generation.²⁾ These findings suggest that the protective effect of US against NO donor-induced neurotoxicity was at least partly a result of its free radical scavenging activity.

It is well known that phenolic compounds isolated from natural products possess antioxidant and free radical scavenging properties.²²⁾ The present data, and those obtained from other neuronal cell types, such as cultured hippocampal cells,⁹⁾ demonstrated that phenolic compounds can inhibit the neuronal damage from the oxidative stress produced by NO generation. Phenolic compounds also have protective effects against oxidative stress-induced cell damage in cultured endothelial cells and ischemia/reperfusion injury by their antioxidant abilities.^{23,24)}

The alkaloid compounds of Uncaria genus have various pharmacological activities, such as vasodilative activity,²⁵⁾ anti-convulsive activity against glutamate-induced convulsion,²⁰⁾ and so forth. In the present study, we demonstrated that not only phenolic fraction but also alkaloid fraction of US have protective abilities against NO-mediated neurotoxicity. We showed previously that an alkaloid fraction of USE suppressed vaso-contraction induced by oxidative stress *in vitro*,²⁵⁾ supporting the notion that alkaloid fraction of US possess free radical scavenging properties.

The neuroprotective effects of traditional Chinese/Japanese (Kampo) medicines besides US were investigated *in vitro*. Watanabe *et al.* reported that the extracts from Toki-shakuyaku-san, a Kampo formula, and its comprised six medicinal plants, Angelicae Radix, Paeoniae Radix, Hoelen, Atractylodis Lanceae Rhizoma,

Alismatis Rhizoma and Cnidii Rhizoma, protected glutamate-induced neuronal damages in CGCs.²⁶⁾ Liu *et al.* reported that Angelicae Radix and Cnidii Rhizoma prevented the methionine sulfoximine-reduced glutamine synthetase activity in mixed CGC-glia cell cultures, and suggested their protective abilities against glutamate-induced neuronal dysfunction.²⁷⁾ Moreover we revealed that the extract from the bark of *Cinnamomum cassia* BLUME, Cinnamomi Cortex, has a protective effect against glutamate-induced neuronal death in CGCs.²⁸⁾

In conclusion, the results of the present study suggest that US has neuroprotective effect against NO-mediated neurotoxicity and that its active components are included in phenolic and alkaloid fractions. This ability of US may have the potential to contribute to protection against NO-related neuronal damage, such as against that occurring in ischemic cerebral diseases.

Acknowledgments

This work was supported by a Health Sciences Research Grant for Comprehensive Research on Aging and Health from the Japanese Ministry of Health, Labour and Welfare.

和文抄録

培養ラット小脳顆粒細胞を用いて、NO donor 誘導神経細胞死に対する釣藤鈎エキスの保護作用を検討した。NO donor には sodium nitroprusside (SNP) と 3-morpholinopyrrolidine (SIN-1) を用い、細胞生存率の評価には MTT 法を用いた。釣藤鈎エキス (10, 30, 100 μ g/ml) は、SNP (30 μ M) 6, 12, 24時間添加による神経細胞死を有意に抑制した。釣藤鈎エキス (10, 30 μ g/ml) は、SIN-1 (300 μ M) 6, 12, 24時間添加による神経細胞死を有意に抑制した。釣藤鈎エキスのフェノール画分は、10, 30 μ g/mlの濃度で SNP (30 μ M, 24時間) 添加による神経細胞死を有意に抑制し、同じく 3, 10 μ g/ml の濃度で SIN-1 (300 μ M, 24時間) 添加による神経細胞死を有意に抑制した。釣藤鈎エキスのアルカロイド画分 (30, 100 μ g/ml) は、SNP (30 μ M, 24時間) および SIN-1 (300 μ M, 24時間) 添加による神経細胞死を有意に抑制した。以上より、釣藤鈎エキスは NO donor によって誘導される神経細胞死に対して保護作用を有し、その活性は釣藤鈎のフェノール画分およびアルカロイド画分にあることが示唆された。

*〒930-0194 富山市杉谷 2630

富山医科薬科大学医学部和漢診療学 嶋田 豊

References

- 1) Bolanos, J.P. and Almeida, A.: Roles of nitric oxide in brain hypoxia-ischemia. *Biochim. Biophys. Acta* **1411**, 415-436, 1999.
- 2) Lipton, S.A. and Rosenberg, P.A.: Excitatory amino acid as a final common pathway for neurologic disorders. *N. Engl. J. Med.* **330**, 613-622, 1994.
- 3) Rothman, S.M. and Olney, J.W.: Glutamate and the pathophysiology of hypoxic-ischemic brain damage. *Ann. Neurol.* **19**, 105-111, 1986.
- 4) Dawson, V.L., Dawson, T.M., London, E.D., Bredt, D.S. and Snyder, S.H.: Nitric oxide mediates glutamate neurotoxicity in primary cortical cultures. *Proc. Natl. Acad. Sci. USA* **88**, 6368-6371, 1991.
- 5) Nomura, Y. and Kitamura, Y.: Inducible nitric oxide synthase in glial cells. *Neurosci. Res.* **18**, 103-107, 1993.
- 6) Lafon-Cazal, M., Pietri, S., Culcasi, M. and Bockaert, J.: NMDA-dependent superoxide production and neurotoxicity. *Nature* **364**, 535-537, 1993.
- 7) Bonfoco, E., Krainc, D., Ankarcrona, M., Nicotera, P. and Lipton, S.A.: Apoptosis and necrosis: two distinct events induced, respectively, by mild and intense insults with *N*-methyl-D-aspartate or nitric oxide/superoxide in cortical cell cultures. *Proc. Natl. Acad. Sci. USA* **92**, 7162-7166, 1995.
- 8) Zhang, J., Dawson, V.L., Dawson, T.M. and Snyder, S.H.: Nitric oxide activation of poly(ADP-ribose) synthetase in neurotoxicity. *Science* **263**, 687-689, 1994.
- 9) Bastianetto, S., Zheng, W.-H. and Quirion, R.: Neuroprotective abilities of resveratrol and other red wine constituents against nitric oxide-related toxicity in cultured hippocampal neurons. *Br. J. Pharmacol.* **131**, 711-720, 2000.
- 10) Lafon-Cazal, M., Culcasi, M., Gaven, F., Pietri, S. and Bockaert, J.: Nitric oxide, superoxide and peroxynitrite: putative mediators of NMDA-induced cell death in cerebellar granule cells. *Neuropharmacology* **32**, 1259-1266, 1993.
- 11) Hogg, N., Darley-Usmar, M., Wilson, M.T. and Moncada, S.: Production of hydroxyl radicals from the simultaneous generation of superoxide and nitric oxide. *Biochem. J.* **281**, 419-424, 1992.
- 12) Shimada, Y., Terasawa, K., Yamamoto, T., Maruyama, I., Saitoh, Y., Kanaki, E. and Takaori, S.: A well-controlled study of Choto-san and placebo in the treatment of vascular dementia. *J. Trad. Med.* **11**, 246-255, 1994.
- 13) Terasawa, K., Shimada, Y., Kita, T., Yamamoto, T., Tosa, H., Tanaka, N., Saito, Y., Kanaki, E., Goto, S., Mizushima, N., Fujioka, M., Takase, S., Seki, H., Kimura, I., Ogawa, T., Nakamura, S., Araki, G., Maruyama, I., Maruyama, Y. and Takaori, S.: Choto-san in the treatment of vascular dementia: a double-blind, placebo-controlled study. *Phytomedicine* **4**, 15-22, 1997.
- 14) Shimada, Y., Goto, H., Kogure, T., Shibahara, N., Kita, T., Itoh, T. and Terasawa, K.: Extract prepared from the hooks and stems of *Uncaria sinensis* prevents glutamate-induced neuronal death in cultured cerebellar granule cells. *J. Trad. Med.* **15**, 141-146, 1998.
- 15) Shimada, Y., Goto, H., Itoh, T., Sakakibara, I., Kubo, M., Sasaki, H. and Terasawa, K.: Evaluation of the protective effects of alkaloids isolated from the hooks and stems of *Uncaria sinensis* on glutamate-induced neuronal death in cultured cerebellar granule cells from rats. *J. Pharm. Pharmacol.* **51**, 715-722, 1999.
- 16) Shimada, Y., Goto, H., Kogure, T., Shibahara, N., Sakakibara, I., Sasaki, H. and Terasawa, K.: Protective effect of phenolic compounds isolated from the hooks and stems of *Uncaria sinensis* on glutamate-induced neuronal death. *Am. J. Chin. Med.* **29**, 173-180, 2001.
- 17) Sakakibara, I., Takahashi, H., Terabayashi, S., Yuzurihara, M., Kubo, M., Higuchi, M., Ishige, A., Komatsu, Y., Maruno, M. and Okada, M.: Chemical and pharmacological evaluations of Chinese crude drug "Gou-teng". *Natural Med.* **52**, 353-359, 1998.
- 18) Sher, P.K. and Hu, S.X.: Increased glutamate uptake and glutamine synthetase activity in neuronal cell cultures surviving chronic hypoxia. *Glia* **3**, 350-357, 1990.
- 19) Liu, J. and Mori, A.: Antioxidant and free radical scavenging activities of *Gastrodia elata* Bl. and *Uncaria rhynchophylla* (Miq.) Jacks. *Neuropharmacology* **31**, 1287-1298, 1992.
- 20) Mimaki, Y., Toshimizu, N., Yamada, K. and Sashida, Y.: Anti-convulsion effects of Choto-san and Chotoko (*Uncaria sinensis* Ramulus) in mice, and identification of the active principles. *Yakugaku Zasshi* **117**, 1011-1021, 1997.
- 21) Hsieh, C.-L., Tang, N.-Y., Chiang, S.-Y., Hsieh, C.-T. and Lin, J.-G.: Anticonvulsive and free radical scavenging actions of two herbs, *Uncaria rhynchophylla* (Miq) Jack and *Gastrodia elata* Bl., in kainic acid-treated rats. *Life Sci.* **65**, 2071-2082, 1999.
- 22) Plumb, G.W., De Pascual-Teresa, S., Santos-Buelga, C., Cheynier, V. and Williamson, G.: Antioxidant properties of catechins and proanthocyanidines: effect of polymerisation, galloylation and glycosylation. *Free Rad. Res.* **29**, 351-358, 1998.
- 23) Law, A., Wu, J., Zeng, L.-H. and Wu, T.-W.: Aortic endothelial cells damaged by a nitric oxide donor and protected by flavonoids. *Life Sci.* **64**, 199-204, 1999.
- 24) Facinó R.M., Carini, M., Aldini, G., Berti, F., Rossoni, G., Bombardelli, E. and Morazzoni, P.: Procyranidines from *Vitis vinifera* seeds protect rabbit heart from ischemia/reperfusion injury: antioxidant intervention and/or iron and copper sequestering ability. *Planta Med.* **62**, 495-502, 1996.
- 25) Goto, H., Sakakibara, I., Shimada, Y., Kasahara, Y. and Terasawa, K.: Vasodilator effect of extract prepared from *Uncaria ramulus* on isolated rat aorta. *Am. J. Chin. Med.* **28**, 197-203, 2000.
- 26) Watanabe, Y., Zhang, X.-Q., Liu, J.-S., Guo, Z., Ohnishi, M. and Shibuya, T.: Protection of glutamate induced neuronal damages in cultured cerebellar granule cells by Chinese herbal medicine, Toki-shakuyaku-san and its comprised six medicinal herbs. *J. Trad. Med.* **12**, 93-101, 1995.
- 27) Liu, J.-S., Watanabe, Y. and Shibuya, T.: Stimulative and protective effects of Chinese medicinal brain metabolic stimulants, Toki, Senkyu and Tanjin, on brain glutamine synthetase activity. *J. Trad. Med.* **12**, 220-228, 1995.
- 28) Shimada, Y., Goto, H., Kogure, T., Kohta, K., Shintani, T., Itoh, T. and Terasawa, K.: Extract prepared from the bark of *Cinnamomum cassia* Blume prevents glutamate-induced neuronal death in cultured cerebellar granule cells. *Phytother. Res.* **14**: 466-468, 2000.