

Inhibitory effects of Cinnamomi Cortex and cinnamaldehyde on oxygen-derived free radical-induced vasocontraction in isolated aorta of spontaneously hypertensive rats

Yuji KASAHARA,^{*a)} Hirozo GOTO,^{b)} Yutaka SHIMADA,^{a)} Nobuyasu SEKIYA,^{a)}
Qiao YANG,^{a)} and Katsutoshi TERASAWA^{a)}

^{a)}Department of Japanese Oriental (Kampo) Medicine, Toyama Medical and Pharmaceutical University,

^{b)}Department of Kampo Diagnostics, Institute of Natural Medicine, Toyama Medical and Pharmaceutical University,
2630 Sugitani, Toyama 930-0194, Japan.

(Received November 5, 2001. Accepted November 16, 2001.)

Abstract

We examined the inhibiting effect of Cinnamomi Cortex extract (CCE) and cinnamaldehyde (CA) against vasocontraction induced by oxygen-derived free radicals produced by the xanthine-xanthine oxidase (XOD) system in the thoracic aortic ring of the spontaneously hypertensive rat (SHR), using the organ bath method *in vitro*.

The vasocontraction induced by xanthine-XOD in the CCE (10^{-4} g/ml) and CA (10^{-4} M) treatment groups were significantly lower than that in the control group. Further, the amounts of thromboxane B₂ (TXB₂) produced in the vasocontractive response in the CCE (10^{-4} g/ml) and CA (10^{-4} M) treatment groups were significantly lower than that in the control group.

For the purpose of examining the mechanism of the inhibiting effect of CA against thromboxane production, the inhibiting effect of CA against the vasocontraction induced by phospholipase A₂ (PLA₂) was examined. The vasocontraction induced by PLA₂ in the CA (10^{-4} M) treatment group was significantly lower than that in the control group. Moreover, the amount of TXB₂ produced by the vasocontractive response in the CA (10^{-4} M) treatment group was significantly lower than that in the control group.

From the above findings, it is suggested that Cinnamomi Cortex is an agent which exerts an inhibitory effect on the vasocontractive factor (TXA₂) *in vitro*.

Key words Cinnamomi Cortex, cinnamaldehyde, vasocontraction, spontaneously hypertensive rats, phospholipase A₂, thromboxane A₂, endothelial dysfunction.

Abbreviations CA, cinnamaldehyde; CCE, Cinnamomi Cortex extract; KB, Keishi-bukuryo-gan (Gui-Zhi-Fu-Ling-Wan), 桂枝茯苓丸; L-NAME, N^G-nitro-L-arginine methyl ester; NO, nitric oxide; PLA₂, phospholipase A₂; SHR, spontaneously hypertensive rat; SOD superoxide dismutase; TXA₂, thromboxane A₂; TXB₂, thromboxane B₂; WKY, Wistar-Kyoto rat; XOD, xanthine oxidase.

Introduction

Vascular endothelial cells produce and secrete various vascular relaxing factors such as nitric oxide (NO),¹⁾ prostaglandin I₂,²⁾ and endothelium-derived hyperpolarizing factor as well as contracting factors such as endothelin and thromboxane A₂ (TXA₂).³⁾ These factors influence each other to create a balance, they maintain a

vascular tonus, and they are involved in blood pressure regulation.¹⁾ There have been many reports concerning the fact that hypertension and vascular endothelial dysfunction are closely related.^{4,5)} Vascular endothelial dysfunction stimulates remodelling of the vascular structure from both a decrease of relaxing factors⁶⁾ and an increase of contracting factors.⁷⁾ Arteriosclerosis progresses, and finally internal organ disorder results.

In our previous study, we have reported on the pro-

*To whom correspondence should be addressed.

protective effect of Keishi-bukuryo-gan (KB; Gui-Zhi-Fu-Ling-Wan) against vascular endothelial disorder in spontaneously hypertensive rats (SHR).⁸⁾ We have found that the contraction induced by xanthine-xanthine oxidase (XOD) is significantly lower in the KB group than in the control group.

The contractive response induced by xanthine-XOD is considered to represent a disordered vascular model caused by arteriosclerosis. The participation against this response of an increase in the production of free radicals in the vascular endothelium and TXA₂, a kind of contracting factor, is reported.⁹⁾

It is reported that Cinnamomi Cortex, the main component of KB, induces endothelial-dependent vascular relaxation¹⁰⁾ and has an improving effect on blood circulation.¹¹⁾ However, the effect of Cinnamomi Cortex on the vasocontractive response has not been confirmed.

In this study, we examine the effect of Cinnamomi Cortex extract (CCE) on the contraction induced by oxygen-derived free radicals produced by the xanthine-XOD system and on TXA₂ production *in vitro*.

Materials and Methods

Preparation of extracts and chemicals: Keihi (桂皮), Cinnamomi Cortex, *Cinnamomum cassia* BLUME, (China), were purchased from Tochimoto Tenkaido (Osaka, Japan). The extract was obtained by boiling these materials in water for 50 minutes and then freeze-drying into a resultant powder. We obtained 3.0g of CCE from 100 g of the raw materials. The CCE powder was then dissolved in distilled water for the experiments. KCl, indomethacin, N^G-nitro-L-arginine methyl ester (L-NAME), superoxide dismutase (SOD) from bovine erythrocytes, dimethyl sulfoxide, xanthine and XOD in suspension were purchased from Wako Pure Chemical Industries (Osaka, Japan). Phospholipase A₂ (PLA₂) from bee venom (*Apis mellifera*) and catalase from bovine liver were purchased from SIGMA-ALDRICH JAPAN K.K. (Tokyo Japan). Trans-cinnamaldehyde was purchased from TOKYO KASEI KOGYO CO., LTD. (Tokyo Japan). Indomethacin and CA were dissolved in dimethyl sulfoxide, and other reagents were dissolved in distilled water and used.

Test animals: SHR and Wistar-Kyoto rats (WKY; 15-20 weeks old, male) were purchased from Sankyo

Labo Service (Tokyo, Japan). Experimental protocols met the "Guidelines for Animal Experimentation" approved by the Japanese Association of Laboratory Animal Science and the Japanese Pharmacological Society.

Contraction experiments: The rats were anesthetized with pentobarbital sodium (50 mg/kg, i.p.) and killed by drawing blood from the heart. A section of the thoracic aorta was carefully cleaned by removing fat and connective tissues, and ring preparations (3 mm wide) were prepared. The rings were mounted on steel hooks in a Magnus chamber (Kishimoto UC-5TD, Kyoto, Japan). One end of the aorta was attached to a force-displacement transducer (Kishimoto UM-203), and its isometric contraction was recorded (Niko Bioscience T-634, Tokyo, Japan). Baths were filled with 7 ml of Krebs solution with the following composition (mM): NaCl 120, KCl 4.7, NaHCO₃ 25.0, KH₂PO₄ 1.2, MgSO₄·7H₂O 1.2, CaCl₂ 2.5, and glucose 10.0. The solution was maintained at 37°C and bubbled continuously with 5% CO₂ - 95% O₂ at pH 7.4. The rings were equilibrated for 40 min at an initial resting tension of 1 g, and then contracted with 60 mM KCl. When contraction reached a steady maximal response, 10⁻⁶ M acetylcholine was added. Then the Krebs solution was replaced every 15 minutes for 60 minutes. Finally, the preparations with the endothelium intact were exposed to L-NAME (10⁻⁴ M) for 60 minutes before contraction.

a) Contraction induced by xanthine-XOD: After it was confirmed that contraction returned to the baseline, each aortic strip was pretreated with CCE (10⁻⁴-10⁻⁶ g/ml); the main component of Cinnamomi Cortex, CA (10⁻⁴ M-10⁻⁶ M); cyclooxygenase inhibitor, indomethacin (10⁻⁵ M); superoxide anion scavenger, SOD (150 U/ml); and hydrogen peroxide catabolic enzyme, catalase (1000 U/ml). At 10 minutes after the addition of the pretreatment drugs, pretreatment with xanthine (10⁻⁴ M) was performed, and 10 minutes after that, XOD (10 mU/ml) was added. Transient contraction was thereby induced. The contraction was expressed as a percentage of 60 mM KCl maximum contraction.

b) Contraction induced by xanthine-XOD with indomethacin: After it was confirmed that contraction returned to the baseline, each aortic strip was pretreated with indomethacin (10⁻⁵ M). At 10 minutes after its addition, each aortic strip was pretreated with CCE (10⁻⁴

g/ml), CA (10^{-4} M) or SOD (150 U/ml). At 10 minutes after the addition of pretreatment drugs, xanthine (10^{-4} M) was added, and XOD (10 mU/ml) was added 10 minutes later. This procedure induced transient contraction. The contraction was expressed as a percentage of the maximum contraction at 60 mM KCl.

c) *Contraction induced by PLA₂*: To examine the effect of CA against TXA₂-induced contraction, PLA₂ (1 U/ml) was administered at 10 minutes after the addition of pretreatment drugs, and transient contraction was induced. The contraction was expressed as a percentage of 60 mM KCl maximum contraction.

Thromboxane B₂ (TXB₂) measurement: In the a) and c) contraction experiments, organ-chamber solutions (2 ml) were collected, respectively, before xanthine pretreatment, 30 minutes after XOD treatment, just before PLA₂ administration, and 30 minutes after PLA₂ administration, and they were used for TXB₂ measurement. TXA₂ is an extremely unstable molecule and rapidly undergoes hydrolytic degradation to stable TXB₂,¹²⁾ which was used as an index of *in vivo* TXA₂ generation. TXB₂ concentration in bath solution was measured by RIA PEG method.¹³⁾

Statistical analysis: Data were presented as mean $\pm$ standard error. Statistical comparisons were done by Mann-Whitney test and repeated measures ANOVA, followed by Fisher's PLSD for multicomparisons, and all values were compared with SHR control values. The level of statistical significance was defined as $p < 0.05$.

Results

Contraction induced by oxygen-derived free radicals produced by the xanthine-XOD system in SHR and WKY

Xanthine (10^{-4} M) plus XOD (10 mU/ml) induced a temporary contraction of the aortic strips with L-NAME. The typical contraction responses are shown in Fig.1A. Contraction of SHR group induced by xanthine-XOD was $73.7 \pm 3.19\%$ ($n = 10$), and that of the WKY group was $11.4 \pm 3.58\%$ ($n = 4$); that of the indomethacin-treated (10^{-5} M) SHR group was $9.32 \pm 3.30\%$ ($n = 10$), that of the SOD-treated (150 U/ml) SHR group was $40.2 \pm 11.01\%$ ($n = 10$), and that of the catalase-treated (1000 U/ml) SHR group was $27.0 \pm 6.25\%$ ($n = 4$; Fig.2). The contractions were significantly greater in the SHR control group than in the WKY group. Both free




Figure 1. Typical tracings of vasocontraction of isolated SHR and WKY aortas after reagent (10^{-5} M indomethacin (IND), 10^{-4} g/ml Cinnamomi Cortex extract (CCE) or 10^{-4} M cinnamaldehyde (CA)) treatment.

- a) Ring contracted with xanthine oxidase (XOD; 10 mU/ml) in the presence of xanthine (Xan; 10^{-4} M).
b) Ring contracted with phospholipase A₂ (PLA₂; 1 U/ml). All aortas had intact endothelium, and had been treated with L-NAME.




Figure 2. Inhibitory effects of reagent (10^{-5} M indomethacin, 150 U/ml SOD or 1000 U/ml catalase) treatment on XOD (10 mU/ml) in the presence of xanthine (10^{-4} M) - induced vasocontraction in isolated SHR and WKY aortas. All aortas had intact endothelium, and had been treated with L-NAME. Contraction was expressed as a percentage of 60 mM KCl maximum contraction. Asterisks indicate significant differences from SHR control group. (* $p < 0.05$; ** $p < 0.01$, mean $\pm$ S.E., $n = 4 \sim 10$)

radical scavengers (SOD and catalase) and cyclooxygenase inhibitor (indomethacin) inhibited the contraction significantly.

Effects of CCE and CA against contraction induced by oxygen-derived free radicals produced by the xanthine - XOD system in SHR

The contraction of the SHR group induced by xanthine-XOD was $73.7 \pm 3.19\%$, that of the CCE-treated (10^{-4} g/ml) SHR group was $25.8 \pm 5.19\%$, and that of the CA-treated (10^{-4} M) SHR group was $31.3 \pm 7.63\%$. Both CCE (10^{-4} g/ml) and CA (10^{-4} M) inhibited the contractions significantly, and both inhibited the con-




Figure 3. Inhibitory effects of reagent (10^{-6} – 10^{-4} g/ml CCE or 10^{-6} – 10^{-4} M CA) treatment on XOD (10 mU/ml) in the presence of xanthine (10^{-4} M) induced vasocontraction in isolated SHR aortas. All aortas had intact endothelium, and had been treated with L-NAME. Contraction was expressed as a percentage of 60 mM KCl maximum contraction. Asterisks indicate significant differences from SHR control group. (* $p < 0.05$; ** $p < 0.01$, mean $\pm$ S.E., $n = 6$ – 10)




Figure 4. Inhibitory effects of the reagent (10^{-4} g/ml CCE, 10^{-4} M CA or 150 U/ml SOD) treatment on XOD (10 mU/ml) in the presence of xanthine (10^{-4} M) - induced vasocontraction in isolated SHR aortas. All aortas had intact endothelium, and had been treated with L-NAME and 10^{-5} M indomethacin. Contraction was expressed as a percentage of 60 mM KCl maximum contraction. (mean $\pm$ S.E., $n = 4$ – 10)

Table I TXB₂ concentration in organ-chamber solution (pg/ml)

Vasocontraction inducer	xanthine (10^{-4} M)-XOD (10 mU/ml)		PLA ₂ (1 U/ml)	
	before	after	before	after
Control(SHR)	9.84 $\pm$ 1.63	91.1 $\pm$ 11.9*	17.6 $\pm$ 7.61	77.6 $\pm$ 14.9*
CCE (10^{-4} g/ml)	12.5 $\pm$ 1.19	34.1 $\pm$ 3.73*#	(–)	(–)
Cinnamaldehyde (10^{-4} M)	9.76 $\pm$ 2.45	47.2 $\pm$ 6.98*#	7.80 $\pm$ 1.96	29.0 $\pm$ 3.94*#

Changes in the concentration of TXB₂ in organ-chamber solution in SHR aortic rings in the vasocontraction inducer after reagent (CCE or CA) treatment. All aortas had intact endothelium, and had been treated with L-NAME.

* $p < 0.05$, vs. before vasocontraction. # $p < 0.05$, vs. SHR control group (mean $\pm$ S.E., $n = 4$ – 7).

tractions in a concentration-dependent manner (each group: $n = 6$ – 10 ; Fig.3).

Effects of CCE and CA against TXB₂ production in the xanthine-XOD-induced contraction

The TXB₂ concentration of the bath solution at 30 minutes after vasocontraction by xanthine-XOD of SHR in the CCE-treated (10^{-4} g/ml) group was significantly lower than that in the control group. The CA (10^{-4} M) group showed a similar result ($n = 6$ – 7 ; Table I).

Contraction induced by oxygen-derived free radicals produced by the xanthine-XOD system in SHR with indomethacin

Contraction of the SHR group with L-NAME and indomethacin induced by xanthine-XOD was $9.4 \pm 3.30\%$ ($n = 10$) that of the CCE-treated (10^{-4} g/ml) SHR group was $6.5 \pm 0.71\%$ ($n = 10$), that of the CA-treated (10^{-4} M) SHR group was $9.2 \pm 2.74\%$ ($n = 6$), and that

of the SOD-treated (150 U/ml) SHR group was $5.3 \pm 0.84\%$ ($n = 4$; Fig.4). In both the indomethacin + CCE and indomethacin + SOD groups, compared with the indomethacin only group, slight inhibition (not significant) of the contraction was seen. But in the indomethacin + CA group, inhibition of the contraction was not observed.

Effect of CA against contraction induced by PLA₂

PLA₂ (1 U/ml) induced a temporary contraction of the aortic strips. The typical contraction responses are shown in Fig.1B. The contraction induced by PLA₂ of the SHR control group was $75.1 \pm 5.03\%$, that of the CA-treated (10^{-4} M) SHR group was $23.2 \pm 8.61\%$, and that of the indomethacin-treated (10^{-5} M) SHR group was $18.0 \pm 6.84\%$. Indomethacin and CA (10^{-4} M) inhibited the vasocontractile response significantly. CA inhibited the vasocontractile response in a concentration-dependent




Figure 5. Inhibitory effects of the reagent (10^{-5} M indomethacin or 10^{-6} - 10^{-4} M CA) treatment on PLA₂ (1 U/ml) - induced vasoconstriction in isolated SHR aortas. All aortas had intact endothelium, and had been treated with L-NAME. Contraction was expressed a percentage of 60 mM KCl maximum contraction. Asterisks indicate significant differences from SHR control group. (* $p < 0.05$, mean $\pm$ S.E., $n = 6 \sim 9$)

manner (each group: $n = 6-9$; Fig.5)

Effect of CA against TXB₂ production in PLA₂-induced contraction

The TXB₂ concentration of the bath solution at 30 minutes after vasoconstriction by PLA₂ of SHR in the CA-treated (10^{-4} M) group was significantly lower than that in the control group ($n = 4-5$; Table I).

Discussion

Cinnamomi Cortex is a clinically important crude drug that is used to improve blood circulation.¹⁴⁾ However, in many respects the mechanisms involved in this improvement are still unclear. One of these mechanisms was reported to be the endothelial-dependent relaxing effect of tannin present in Cinnamomi Cortex.¹⁰⁾ In this study, as the influence of NO was excluded by pretreatment with L-NAME, other mechanisms were studied.

It is known that vessel injury by atherosclerosis produces free radicals. Such free radicals have already been produced by combining xanthine and XOD *in vitro*.⁹⁾ When one molecule of uric acid is formed into one molecule of xanthine by XOD, one molecule of superoxide radical is also formed. It is reported that, when this superoxide radical acts on blood vessels, it causes an increase in TXA₂ in the blood vessels and, in turn, their contraction.⁹⁾ These phenomena were also observed in the present study (Fig.1 and 2, Table I). The contraction

induced by xanthine and XOD was significantly inhibited by pretreatment with radical scavengers (SOD and catalase; Fig.2). From these findings, it is considered that this contraction response was induced by free radicals.

The xanthine-XOD-induced contraction was inhibited by pretreatment with CCE. The increase of TXB₂ in the organ bath was suppressed by CCE compared to the control group (Fig.3, Table I).

TXA₂ is produced not only in platelets¹²⁾ but also in the endothelium and smooth muscle of vessels.¹⁵⁾ In this study, TXB₂ was considered to originate from blood vessels¹⁶⁾ and, as a result, TXA₂ production was thought to be inhibited.

Further, cinnamaldehyde, a main component of Cinnamomi Cortex, was also studied. The contraction induced by xanthine and XOD was inhibited by pretreatment with cinnamaldehyde in the same way as with CCE (Fig.3). This suggested that cinnamaldehyde was one of the main components in the inhibiting effect for xanthine and XOD contraction. Moreover, the mechanism was considered to be related to the production of TXA₂.

The contraction induced by xanthine-XOD was inhibited by pretreatment with indomethacin (Fig.2, Table I). It is thought that superoxide radicals stimulate the arachidonate cascade, and thus TXA₂ is produced. This agrees with the past report that free radical-induced contraction was inhibited by pretreatment with an inhibitor of PLA₂, the first enzyme of the arachidonate cascade.¹⁷⁾ We considered that these inhibitory effects of CCE and CA on the xanthine-XOD-induced contraction are the free radical scavenging effect, or the inhibitory effect, of the enzyme activity of a part of the arachidonate cascade. So we examined the free radical scavenging effect of CCE and CA.

The xanthine-XOD-induced contraction with indomethacin was smaller than the contraction without indomethacin (Fig.2), and was even more inhibited by pretreatment with CCE and SOD. It is suggested that this additional effect is the radical scavenging effect of SOD. Also, it is reported that the tannin present in Cinnamomi Cortex possesses a radical scavenging effect *in vitro*.¹⁰⁾ As for the additional effect by CCE, it is also thought to be a radical scavenging effect. On the other hand, the contraction was not inhibited by pretreatment with CA (Fig.4), and it was considered that CA does not exert a radical scavenging effect.

To determine the effect of CA on the system of TXA₂ production, we studied its effect on PLA₂-induced contraction. We found that pretreatment with CA inhibited PLA₂-induced contraction in a gradual, dose-dependent manner (Fig.5). It has been reported that CA has an inhibitory effect on thromboxane production that results from the inhibitory effect of the arachidonic acid release by the suppressive effect of PLA₂ activity on platelet aggregation *in vitro*.¹⁸⁾ Thus, it is possible that CA inhibits the PLA₂ activity on vasocontraction *in vitro*. It is suggested that this contraction inhibitory effect of CA is a result of its inhibitory effect on PLA₂ activity.

In earlier studies, it has been reported that drugs exerting a suppressive effect on platelet aggregation were paeonol from Moutan Radix Cortex¹⁹⁾ and trichosanic acid from Trichosanthis Semen.²⁰⁾ But there has been no report on the effect on TXA₂ production in injured vessels.

TXA₂ has strong activity to contract vessels, so it is closely related to hypertension. KB has been reported to decrease blood pressure,⁸⁾ and this is thought to be affected by the suppression of TXA₂ production. Thus, from the present results it is suggested that CCE works to decrease blood pressure and improve related symptoms.

Further, the effects of Toki-sigyakuka-goshuyushokyo-to on peripheral circulation²¹⁾ and of Sai-rei-to on TXA₂ metabolism in nephrotic rat have been reported.²²⁾ It was suggested that these effects were concerned with the suppressive effect of TXA₂ production by CA.

It has also been reported that TXA₂ has proliferating and hypertrophic effects on smooth muscle,⁷⁾ and the accumulation of the expression of mRNA of the thromboxane receptor has been recognized in the arteriosclerotic lesions of human, even in the absence of coronary disease.²³⁾ In addition, it was reported that the serum TXA₂ concentration of a youth with genetically inherited coronary disease diathesis was elevated.²⁴⁾ The accumulated evidence strongly suggests that TXA₂ is involved in arteriosclerotic development. Taken together then, the possibility that Cinnamomi Cortex has an anti-atherosclerotic effect from its suppressive effect on TXA₂ production cannot be denied.

In our past study, we have reported that tannin in Cinnamomi Cortex induced endothelial-dependent vas-

cular relaxation *in vitro*.¹⁰⁾ When that finding and our current results are examined together, we are convincingly led to the assumption that Cinnamomi Cortex is an agent with the versatility of possessing a potent effect that manifests itself both as a vascular relaxing factor and as suppressive activity against the vasoconstrictive factor (TXA₂) *in vitro*.

Acknowledgments

This study was supported by a grant-in-aid for Funds for Comprehensive Research on Aging and Health from the Japanese Ministry of Health, Labor and Welfare.

和文抄録

自然発症高血圧ラット (SHR) 摘出胸部大動脈におけるキサンチン-キサンチンオキシダーゼ (XOD) 誘発血管収縮反応に対する桂皮エキスおよびケイヒアルデヒド (CA) の収縮抑制作用についてオルガンバス法を用いて検討した。

キサンチン-XOD 誘発血管収縮反応は, SHR 対照群と比較して, 桂皮エキス (10⁻⁴ g/ml) 前処置群, CA (10⁻⁴ M) 前処置群で, 有意に抑制されていた。キサンチン-XOD 収縮反応時のトロンボキサン B₂ (TXB₂) 産生量は, SHR 対照群と比較して, 桂皮エキス (10⁻⁴ g/ml) 群, CA (10⁻⁴ M) 群で, 有意に抑制されていた。

CA の TX 産生抑制の機序を検討するため, フォスホリパーゼ A₂ (PLA₂) 誘発血管収縮反応に対する収縮抑制作用について検討したところ, PLA₂誘発血管収縮反応は SHR 対照群と比較して, CA (10⁻⁴ M) 群で, 有意に抑制されていた。PLA₂ 収縮反応時の TXB₂ 産生量は, SHR 対照群と比較して CA (10⁻⁴ M) 群で, 有意に抑制されていた。

以上のことから, 桂皮は, 血管収縮因子である TXA₂ 抑制作用を持つ生薬である可能性が示唆された。

*〒930-0194 富山市杉谷 2630

富山医科薬科大学医学部和漢診療学講座 笠原裕司

References

- 1) Vanhoutte, P.M.: Endothelium and control of vascular function. *Hypertension* **13**, 658-667, 1989.
- 2) Moncada, S., Herman, A.G., Higgs, E.A. and Vane JR.: Differential formation of prostacyclin (PGX or PGI₂) by layers of

- the arterial wall. An explanation for the anti-thrombotic properties of vascular endothelium. *Thrombosis Research* **11**, 323-344, 1977.
- 3) Lüscher, T.F.: The endothelium target and promoter of hypertension? *Hypertension* **15**, 482-485, 1990.
 - 4) McGuire, P.G. and Twietmyer, T.A.: Aortic endothelial junctions in developing hypertension. *Hypertension* **7**, 483-490, 1985.
 - 5) Ferrante, F., Ciriaco, E., Abbate, F., Laura, R. and Amenta, F.: Effect of long term isradipine treatment on the morphology of the endothelium in the aorta of spontaneously hypertensive rats. *Clin. Exper. Hypertension* **16**, 865-880, 1994.
 - 6) Leyen, H.E., Gibbons, G.H., Morishita, R., Lewis, N.P., Zhang, L., Nakajima, M., Kaneda, Y., Cooke, J.P. and Dzau, V.J.: Gene therapy inhibiting neointimal vascular lesion. In vivo transfer of endothelial cell nitric oxide synthase gene. *Proc. Natl. Acad. Sci. USA* **92**, 1137-1141, 1995.
 - 7) Dzau, V.J.: The role of mechanical and hormonal factors in growth regulation of vascular smooth muscle and cardiac myocytes. *Curr. Opin. Nephrol. Hypertens.* **2**, 27-32, 1993.
 - 8) Kasahara, Y., Goto, H., Shimada, Y., Sekiya, N., Yang, Q., and Terasawa, K.: Effects of Keishi-bukuryo-gan (Gui-Zhi-Fu-Ling-Wan) on endothelial function in spontaneously hypertensive rats. *J. Trad. Med.* **18**, 113-118, 2001.
 - 9) Hibino, M., Okumura, K., Iwama, Y., Mokuno, S., Osanai, H., Matsui, H., Toki, Y. and Ito, T.: Oxygen-Derived Free Radical-Induced Vasoconstriction by Thromboxane A₂ in Aorta of the Spontaneously Hypertensive Rat. *J. Cardiovasc. Pharmacol.* **33**, 605-610, 1999.
 - 10) Tanikawa, K., Goto, H., Nakamura, N., Tanaka, N., Hattori, M., Itoh, T. and Terasawa, K.: Endothelium-dependent vasodilator effect of tannin extract from Cinnamomi Cortex on isolated rat aorta. *J. Trad. Med.* **16**, 45-50, 1999.
 - 11) Harada, M. and Yano, S.: Pharmacological studies on Chinese cinnamon. II. Effects of cinnamaldehyde on the cardiovascular and digestive systems. *Chem. Pharm. Bull.* **23**, 941-947, 1975.
 - 12) Oates, J.A., FitzGerald, G.A., Branch, R.A., Jackson, E.K., Knapp, H.R. and Roberts, L.J.: II. Clinical implications of prostaglandin and thromboxane A₂ formation. *N. Engl. J. Med.* **319**, 689-698, 1988.
 - 13) Morris, H.G., Sherman, N.A., Shepperdson, F.T.: Variables associated with radioimmunoassay of prostaglandins in plasma. *Prostaglandins* **21**, 771-788, 1981.
 - 14) Terasawa, K.: *Kampo Japanese-Oriental Medicine*. pp. 43-46, pp. 200-201, K.K.Standard McIntyre, Tokyo, Japan, 1993.
 - 15) Sachinidis, A., Flesch, M., Ko, Y., Schrör, K., Böhm, M., Düsing, R. and Vetter, H.: Thromboxane A₂ and vascular smooth muscle cell proliferation. *Hypertension* **26**, 771-780, 1995.
 - 16) Uehara, Y., Tobian, L., Iwai, J., Ishii, M. and Sugimoto, T.: Alterations of vascular prostacyclin and thromboxane A₂ in Dahl genetical strain susceptible to salt-induced hypertension. *Prostaglandins* **33**, 727-738, 1987.
 - 17) Peters, S.L.M., Mathy, M.-J., Pfaffendorf, M. and van Zwieten, P.A.: Reactive oxygen species-induced aortic vasoconstriction and deterioration of functional integrity. *Naunyn-Schmiedeberg's Arch Pharmacol.* **361**, 127-133, 2000.
 - 18) Takenaga, M., Hirai, A., Terano, T., Tamura, Y., Kitagawa, H. and Yoshida, S.: In vitro effect of cinnamic aldehyde, a main component of Cinnamomi Cortex, on human platelet aggregation and arachidonic acid metabolism. *J. Pharmacobio-Dyn.* **10**, 201-208, 1987.
 - 19) Hirai, A., Terano, T., Hamazaki, T., Sajiki, J., Saito, H., Tahara, K., Tamura, Y. and Kumagai, A.: Studies on the mechanism of antiaggregatory effect of Moutan Cortex. *Thrombosis Research* **31**, 29-40, 1983.
 - 20) Takenaga, M., Hirai, A., Terano, T., Tamura, Y., Kitagawa, H. and Yoshida, S.: In vitro effect of Trichosanin acid, a major component of Trichosanthes Japonica on platelet aggregation and arachidonic acid metabolism in human platelets. *Prostaglandins, Leukotrienes and Essential Fatty Acids* **31**, 65-72, 1988.
 - 21) Kanai, S., Okano, H., and Abe, H.: Efficacy of Toki-shigyakukagoshiyuu-syokyo-to (Danggui-Sini-Jia-Wuzhuyu-Shengjiang-Tang) on Peripheral Circulation in Autonomic Disorders. *Am. J. Chin. Med.* **25**, 69-78, 1997.
 - 22) Suzuki, J. and Nagasawa, K.: Effect of Sai-rei-to on nephrotic rat - examination from the TXA₂ metabolism. (Japanese) *Nihon Toyo Igaku Zasshi (J. J. Orient. Med.)* **45**, suppl., 128, 1997.
 - 23) Azakami, S.: Effect of thromboxane A₂ in human atherosclerotic lesion. (Japanese) *Kurume Igakukai Zasshi* **57**, 1261-1270, 1994.
 - 24) Akimova, E.V.: Prostacyclin and thromboxane A₂ levels in children and adolescents with an inherited predisposition to coronary heart disease: a family study. *Coronary Artery Disease* **5**, 761-765, 1994.